

See FDA package inserts for more information, including more detailed safety information. Ask your doctor if one of these options is right for you.

Medication	Cautions/Warnings	Side Effects	Dosage	Use	Availability (check insurance)
Bupropion SR 150	Not for use if you: * Use monoamine oxidase (MAO) inhibitor * Use bupropion in any other form * Have a history of seizures * Have a history of eating disorders FDA Boxed Warning: See the FDA Web Site for more information	* Insomnia * Dry mouth	* Days 1-3: 150 mg each morning * Days 4–end: 150 mg twice daily	Start 1-2 weeks before quit date; use 2 to 6 months	Prescription Only: * Generic * Zyban * Wellbutrin SR
Nicotine Gum (2 mg or 4 mg)	* Caution with dentures * Do not eat or drink 15 minutes before or during use	* Mouth soreness * Stomach ache	* 1 piece every 1 to 2 hours * 6-15 pieces per day * If ≤ 24 cigs: 2 mg * If ≥ 25 cigs/day: 4 mg	Up to 12 weeks or as needed	OTC Only: * Generic * Nicorette
Nicotine Inhaler	* May irritate mouth/throat at first (but improves with use)	* Local irritation of mouth & throat	* 6-16 cartridges/day * Inhale 80 times/cartridge * May save partially-used cartridge for next day	Up to 6 months; taper at end	Prescription Only: * Nicotrol inhaler
Nicotine Lozenge (2 mg or 4 mg)	* Do not eat or drink 15 minutes before or during use * One lozenge at a time * Limit 20 in 24 hours	* Hiccups * Cough * Heartburn	* If smoke ≥ 30 minutes after waking: 2 mg * If smoke ≤ 30 minutes after waking: 4 mg * Weeks 1-6: 1 every 1-2 hrs * Wks 7-9: 1 every 2-4 hrs * Wks 10-12: 1 every 4-8 hrs	3-6 months	OTC Only: * Generic * Commit
Nicotine Nasal Spray	* Not for patients with asthma * May irritate nose (improves over time) * May cause dependence	* Nasal irritation	* 1 “dose” = 1 squirt per nostril * 1 to 2 doses per hour * 8 to 40 doses per day * Do NOT inhale	3-6 months; taper at end	Prescription Only: * Nicotrol NS
Nicotine Patch (7 mg, 14 mg or 21 mg)	* Do not use if you have severe eczema or psoriasis	* Local skin reaction * Insomnia	* One patch per day * If ≥ 10 cigs/day: 21 mg 4 wks, 14 mg 2-4 wks, 7 mg 2-4 wks	8-12 weeks	OTC or prescription: * Generic * Nicoderm CQ * Nicotrol
Varenicline	Use with caution in patients: * With significant renal impairment * With serious psychiatric illness * Undergoing dialysis FDA Boxed Warning: See the FDA Web Site for more information	* Nausea * Insomnia * Abnormal, vivid or strange dreams	* Days 1-3: 0.5 mg every morning * Days 4-7: 0.5 mg twice daily * Day 8–end: 1 mg twice daily	Start 1 week before quit date; use 3-6 months	Prescription only: * Chantix
Combinations: 1) Patch + bupropion 2) Patch + gum 3) Patch + [lozenge or inhaler]	* Only patch + bupropion is currently FDA-approved. * Follow instructions for individual medications.	See individual medications above.	See individual medications above.	See above.	See above.